

The amino terminal end determines the stability and assembling capacity of eukaryotic ribosomal stalk proteins P1 and P2

Hendricka Camargo, Gretel Nusspaumer, David Abia, Verónica Briceño, Miguel Remacha and Juan P. G. Ballesta*

Centro de Biología Molecular Severo Ochoa, Consejo Superior de Investigaciones Científicas and Universidad Autónoma de Madrid, Cantoblanco, 28049 Madrid, Spain

Received August 11, 2010; Revised December 10, 2010; Accepted December 28, 2010

ABSTRACT

The eukaryotic ribosomal proteins P1 and P2 bind to protein P0 through their N-terminal domain to form the essential ribosomal stalk. A mutational analysis points to amino acids at positions 2 and 3 as determinants for the drastic difference of *Saccharomyces cerevisiae* P1 and P2 half-life, and suggest different degradation mechanisms for each protein type. Moreover, the capacity to form P1/P2 heterodimers is drastically affected by mutations in the P2 β four initial amino acids, while these mutations have no effect on P1 β . Binding of P2 β and, to a lesser extent, P1 β to the ribosome is also seriously affected showing the high relevance of the amino acids in the first turn of the NTD α -helix 1 for the stalk assembly. The negative effect of some mutations on ribosome binding can be reversed by the presence of the second P1/P2 couple in the ribosome, indicating a stabilizing structural influence between the two heterodimers. Unexpectedly, some mutations totally abolish heterodimer formation but allow significant ribosome binding and, therefore, a previous P1 and P2 association seems not to be an absolute requirement for stalk assembly. Homology modeling of the protein complexes suggests that the mutated residues can affect the overall protein conformation.

INTRODUCTION

The stalk is a lateral protuberance of the large ribosomal subunit involved in the interaction and function of several

soluble factors during translation (1). The ribosomal stalk is formed by a complex of a ribosomal RNA (rRNA)-binding protein, the prokaryotic L10 and the eukaryotic P0, and a set of acidic proteins called L7/L12 in prokaryotes and P1/P2 in eukaryotes. Protein L7/L12 binds as a homodimer to L10 forming either a pentameric L10-(L7/L12)₂ or a heptameric L10-(L7/L12)₃ complex depending on the organism (2,3). In higher eukaryotes there are two acidic protein families, P1 and P2, which interact with protein P0 as P1/P2 heterodimers to form a P0-(P1/P2)₂ pentamer (4,5). *Saccharomyces cerevisiae* contain two different forms in each acidic protein family, α and β , which are found in the ribosome also forming a P0-(P1 α /P2 β)(P1 β /P2 α) pentamer (6).

A notable dynamism is one of most significant features of the eukaryotic stalk. Thus, in contrast to the stability of the prokaryotic L10-L7/L12 interaction (7), the eukaryotic complex is easily disassembled (8). Moreover, the ribosome-bound P1 and P2 proteins are exchanged with the free acidic proteins present in a large cytoplasmic pool (9–11) and this process, which implies changes in the P1/P2 affinity for the ribosome, is increased during protein synthesis (12). This stalk dynamism results in the presence of diverse ribosome subpopulations containing different amounts of P1/P2 proteins (13,14). The absence of P1/P2 reduces the ribosome translating efficiency to a different extent depending on the translated messenger RNAs (mRNAs), and some proteins are expressed at a higher rate by the defective ribosomes (15). Therefore, the overall cellular pattern of protein expression can be defined, among other mechanisms, by the relative proportion of each ribosome subpopulation, which is apparently determined by the metabolic activity of the cell.

To fully understand the proposed stalk-dependent regulation at the molecular level, it is indispensable to know

*To whom correspondence should be addressed. Tel: +34 911964505; Fax: +34 911964420; Email: jpgballesta@cbm.uam.es
Correspondence may also be addressed to Miguel Remacha. Tel: +34 911964506; Fax: +34 911964420; Email: mremacha@cbm.uam.es
Present address:

Gretel Nusspaumer, Pediatric Immunology Laboratory, Department of Biomedicine, UKBB and University of Basel, Mattenstrasse 28, CH-4058 Basel, Switzerland.

the mechanism involved in the assembly and disassembly process and how it can be controlled by the cell. In this regard, there are at least two important aspects to be considered. One of them concerns the interaction dynamics between P1, P2 and P0, while the other is related to the size and control of the P1/P2 cytoplasmic pool, which must affect the equilibrium between the different ribosome subpopulations. Regarding the interaction between the stalk components, the regions determining the binding site have been characterized in both P0 (16–18) and in P1/P2 (19), but a high-resolution structure of the eukaryotic complex, necessary to fully understand the complex formation, is unfortunately missing. The crystal structure of the archaeal ribosomal stalk core structure recently reported (20) can be helpful in understanding the equivalent eukaryotic ribosomal domain, considering the significant similarity of their components. Similarly, the newly reported nuclear magnetic resonance (NMR) structure of a mammalian protein P2 homodimer (21) is an important step toward the resolution of the really biologically important P1/P2 heterodimer.

On the other hand, information available on size and control of the cytoplasmic pool of the P1 and P2 proteins in eukaryotic cells is rather scarce. There are data pointing to an effect at the transcriptional level (22). Moreover, there is a notable difference in the stability of the two acidic protein types, which may have a role in determining their relative proportion in the cytoplasm (23). Thus, while the half-life of P2 proteins is longer than 5 h, that of the P1 proteins is shorter than 15 min; nevertheless, the association of both protein types forming a P1/P2 couple protects P1 from degradation (23,24). This association is therefore essential for the existence of a pool of P1 proteins, and it is also an important fact in understanding the stalk assembly.

We previously reported that the different degradation sensitivity of both acidic protein types, P1 and P2, is determined by the first five amino acids of the protein sequence but not by the N-terminal acetylation (23). In this report, we have analyzed in more detail the degradation determinants at the N-terminal of *S. cerevisiae* P2 β and P1 β proteins by replacing each one of the first four residues in protein P2 β by those present in the same position in protein P1 β and *vice versa*. At the same time, we have checked the effects of mutations in this region on P1–P2 heterodimer formation as well as in the binding of the proteins to the ribosome. Our results, in addition to identifying more precisely the degradation determinants, have shown that modifications at the amino end of the proteins seriously affect their capacity to form P1/P2 couples as well as the protein affinity for P0.

MATERIALS AND METHODS

Strains and growth conditions

Saccharomyces cerevisiae D5 (MAT α ; *leu2-3*, *trp1-1*, *ura3-1*, *ade2-1*, *his3-11,15*, *can 1-100*, *RPP2 β ::HIS3*) and D56 (MAT α ; *leu2-3*, *trp1-1*, *ura3-1*, *ade2-1*, *his3-11,15*, *can 1-100*, 3 *RPP1 β ::TRP1*, *RPP2 β ::HIS*) were derived from

S. cerevisiae W3031b as previously described (24,25). *Saccharomyces cerevisiae* MaV203 (MAT α , *leu2-3,112*, *trp1-901*, *his3 Δ 200*, *ade2-101*, *gal4 Δ* , *gal80 Δ* , SPAL10::URA3, GAL1::lacZ, HIS3_{UAS} GAL1::HIS3@LYS2, *can1^R*, *cyh2^R*) (Invitrogen) was used as reporter host strain in the two-hybrid assay. Yeasts were grown in either rich YEP or minimal SC media supplemented with 2% glucose.

Escherichia coli XL1-Blue was used for plasmid manipulations and was grown in LB medium.

Plasmids

Constructs carrying proteins P1 β and P2 β mutations. The plasmids carrying protein P1 β F2 and P1 β 5ntP2 β have been previously described in refs. (26 and 23), respectively. The remaining constructs were obtained following a similar strategy (23) detailed in the Supplementary Data using oligonucleotides in Table S1.

Two-hybrid constructs. Polymerase chain reaction (PCR) DNA fragments were obtained from the cloned *RPP2B* and *PRP1B* and inserted into plasmids pBDC(*TRP1*) or pADC(*LEU2*) as previously reported (27) and is detailed in the Supplementary Data.

Genetic manipulations and recombinant DNA techniques were carried out following standard protocols (28). PCR reactions were performed with Ex Taq^{RT} polymerase (TaKaRa) and custom-made oligonucleotides (Isogen) according to re. 29.

Cell fractionation and ribosome preparation

Cell extracts and washed ribosomes from *S. cerevisiae* cells were prepared as previously described (19)

Protein analysis

Proteins were analyzed either by 15% sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS–PAGE) or by isoelectrofocusing in 5% polyacrylamide gels using a 2.0–5.0 pH range (30) and detected either by western blotting using a specific anti-P2 β or anti-P1 β monoclonal antibodies (31) or by silver staining.

Protein N-terminal sequences have been obtained by Edman degradation at the Servicio de Proteómica in the Centro de Investigaciones Biológicas, Madrid.

Analysis of protein interactions in a two-hybrid assay

Two-hybrid yeast interaction assays were performed in the *S. cerevisiae* Mav203 strain, which carries the *URA3* gene under control of the GAL4 promoter (32). Pairs of plasmids expressing the activation and binding domain from GAL4 fused to the carboxyl end of the proteins to be tested (27) were co-transformed into the yeast cells and the transformants were selected on solid synthetic minimal medium free of tryptophan and leucine (SC-Trp-Leu). Protein interactions were analyzed in SC-Ura solid medium by plating dilutions of each transformed strain.

Affinity chromatography

Cell supernatant (S100 fraction) and Ni^{2+} -NTA resin (Invitrogen) were mixed in a ratio of 1 mg of extract per 35 μl of resin, previously equilibrated in LY buffer (10 mM Tris-HCl, pH 7.5, 20 mM NaCl, 50 mM KCl, 10 mM MgCl_2). The sample was incubated for 1 h. at 4°C with continuous shaking and then the mixture was loaded in a column (10 cm \times 1 cm), and the flow through fraction was collected. The column was subsequently washed first with 10 vol of buffer LY, then with 10 vol of 10 mM imidazol in buffer LY. Finally, the resin-bound proteins were eluted with 4 vol of 100 mM imidazol in the same buffer and the eluted fraction was collected.

Protein complex structure modeling

Models of *S. cerevisiae* P1/P2 heterodimers and the stalk pentamer have been generated by MODELLER 9.8 (33) using human NTD-P2 (2W1O) (21), and archaeal stalk complex (3A1Y) (20) as templates.

Molecular dynamics simulations were run in NAMD (34) for complexes fully solvated with explicit water molecules, during 20 ns at 298 K under constant pressure and periodic boundary conditions. Further details are available as Supplementary Data.

RESULTS

Residues at positions 2 and 3 in the amino acid sequence are major determinants for degradation sensitivity in proteins P1 β and P2 β

One significant difference compared to prokaryotes in the structure of the NH_2 -terminal end of eukaryotic acidic P1 and P2 proteins is acetylation, which takes place at Ser 2 after removal of the starting methionine only in the P1 proteins (35). Amino-terminal acetylation has been considered as a degradation signal in some proteins (36,37), and, therefore, the effect of modifications in Ser 2 and neighboring residues on the sensitivity to degradation in

protein P1 β was explored. Proteins carrying a mutation in position 2, P1 β F2, and a double mutation in positions 2 and 4, P1 β K2L4, as well as a P1 β derivative with the five first amino acids substituted by the equivalent sequence from P2 β , P1 β 5ntP2 β , were constructed (Table 1). The mutated genes were expressed in *S. cerevisiae* D456 which lacks the wild-type P1 β and the two P2 proteins (15). In this strain, the expressed mutated P1 β proteins do not have their P2 partner and then are unable to bind to the ribosome. They rest unprotected in the cytoplasm and accessible to cellular proteolytic activities (23). As shown in Figure 1A, a western blot analysis of transformed D456 total cell extracts showed that mutation of S2F resulted in an increase of the amount of P1 β to the parental W303 level while the wild-type protein was fully degraded in the same conditions. Additional mutations (P1 β K2L4 and P1 β 5ntP2 β) did not cause a further P1 β increase in the cell extracts (Figure 1A).

Considering the previous results, it could be expected that introducing a Ser at position 2 in P2 proteins destabilizes these highly stable proteins. However, preliminary experiments showed that a Ser at position 2 or/and position 4 in protein P2 β had only a minor effect on the protein stability (data not shown, see below). Therefore, a more extensive analysis was carried out basically by replacing each one of the first four residues in protein P2 β by those present in the same position in protein P1 β (Table 1). The mutant proteins were expressed in *S. cerevisiae* D567, which lacks P2 β and both P1 proteins (15). In this strain, the expressed P2 β protein lacks the corresponding P1 partner and remains free in the cytoplasm. Estimation of the protein in the extracts confirmed that the amount of P2 β is practically unaffected by the introduction of a Ser at position 2 (protein P2 β S2) and is somewhat decreased by an additional Ser at position 4 (protein P2 β S2S4) (Figure 1A). In contrast, the protein is absent in the extracts from all mutants carrying an Asp in position 3 (P2 β D3, P2 β S2D3, P2 β D3S4 and P2 β S2D3S4).

Table 1. Mutations used in this work

Protein	Mutation	N-terminal sequence		
		Encoded ^a	Expected ^b	Sequenced ^c
P1 β	None	MSDSI	Ac-SDSI	Blocked
P1 β F2	S2F	MFDSI	MFDSI	MFDS
P1 β K2L4	S2K,S4L	MKDLI	MKDLI	N/D
P1 β 5ntP2 β	S2K,D3Y,S4L,I4A	MKYLA	MKYLA	N/D
P2 β	None	MKYLA	MKYLA	MKYL
P2 β S2 ^d	K2S	MSYLA	Ac-SYLA	MSYL, SYLA, Blocked
P2 β S2S4	K2S,L4S	MSYSA	Ac-SYSA	Blocked
P2 β S2D3S4	K2S,Y3D,L4S	MSDSA	Ac-SDSA	N/D
P2 β S2D3	K2S,Y3D	MSDLA	Ac-SDLA	N/D
P2 β D3	Y3D	MKDLA	MKDLA	MKDL
P2 β D3S4	Y3D,L4S	MKDSA	Ac-KDSA	N/D

^aSequence derived from gene nucleotide sequence. Mutated positions are in bold.

^bSequence expected from N-terminal processing rules.

^cSequence determined by Edman degradation in proteins able to bind to ribosomes (Figure 2).

^dSequencing has shown the presence of an N-terminal blocked form in minor amounts while two unblocked forms (MSYL, SYLA) are present in similar amounts (Figure 2).




Figure 1. Estimation of soluble protein P1 β and P2 β mutants in *S. cerevisiae*. Total extracts from *S. cerevisiae* D456 (A) or D567 (B) transformed with plasmid expressing the indicated proteins were resolved by SDS-PAGE and the proteins detected by immunoblotting using a specific monoclonal antibody to either protein. Extracts from the parental W303 strain were used as control.

Effect of N-terminal mutations on protein binding to the ribosome

The P1 β and P2 β mutant series were expressed in *S. cerevisiae* D56, which lacks proteins P1 β and P2 β but contains their respective protein partner, P2 α and P1 α (24), and the ribosome bound stalk components were analyzed by isoelectrofocusing (30). The S2F and S2K,S4L mutations in P1 β F2 and P1 β K2L4 had no detectable effect on the amount of ribosome-bound protein, and it had previously been found that a single S4R mutation did not affect binding to the ribosome either (38). In contrast, protein P1 β 5ntP2 β was not detected in the particles (Figure 2A). In the P2 β series, introduction of a Ser at position 2 resulted in a duplication of the mutated protein bands, which correspond to different N-terminal processed forms of the protein (Table 1 and Supplementary Data). The K2S mutation had a limited effect on the protein ribosome-binding capacity. In contrast, the protein in the ribosome was notably affected by mutating position 3 or by a double mutation in two out of the three analyzed amino acids, and was totally abolished by changing the three of them simultaneously (Figure 2B and Supplementary Figure S1). It must be noted that the bound mutant proteins are functional since they stimulate strain D56 growth (Supplementary Data, Table S2).

The P2 β mutants were also expressed in *S. cerevisiae* D5, which only lacks their parental wild-type protein. In this case, after transformation the cells will contain a full complement of stalk components and it will be possible to test whether this fact somehow affects the binding of the mutated proteins to the ribosomes. The ribosomes from both transformed strains were analyzed by SDS-PAGE and the proteins estimated by immunoblotting (Figure 2C). It was clear that the presence of a complete stalk favored the binding of proteins P2 β S2S4, P2 β S2D3 and P2 β D3S4, which were present in higher amounts in ribosomes from strain D5. In contrast, the triple mutant protein was not found in either strain.




Figure 2. Detection of ribosome-bound protein P1 β and P2 β mutants. (A and B) Purified ribosomes from *S. cerevisiae* D56 expressing either the P1 β (A) or the P2 β (B) mutants were resolved by isoelectrofocusing and the proteins detected by silver staining. The P proteins usually appear as a double band, which correspond to the phosphorylated (lower band) and non-phosphorylated (upper band) forms. The mutated proteins are marked by an asterisk in the stained gel. (C) Ribosomes purified from strains D5 and D56 transformed with the P2 β mutants were resolved by SDS-PAGE and the proteins detected by immunoblotting using a specific anti-P2 β monoclonal antibody.

The formation of P1 α /P2 β heterodimers is affected by the N-terminal mutations

The capacity of the P1 and P2 proteins to form P1/P2 heterodimers, preferentially P1 α /P2 β and P1 β /P2 α in the case of *S. cerevisiae* (39), suggested that the formation of these associations is a requirement for binding to the ribosome. Since some of our mutant proteins are either totally or partially affected in ribosome binding we checked whether their ribosome binding defect parallels their capacity to associate with the partner P1 α protein. Thus, we tested the interaction of the P1 β and P2 β mutant series with proteins P2 α and P1 α , respectively, using the two-hybrid technique. Either the tests were carried out with the mutant proteins fused to the GAL4 binding




Figure 3. Interaction of P1β mutants with protein P2α (upper panel) and P2β mutants with protein P1α (lower panel) estimated by the two-hybrid test. *S. cerevisiae* MaV203 was co-transformed with the construct expressing the different mutants fused to the GAL4 binding domain and the plasmid expressing the corresponding wild-type protein fused to the GAL4 activating domain. Serial dilutions of the transformed cells were plated on SC medium lacking uracil (left panels). The extent of growth in these conditions is proportional to the interaction between the tested proteins. As a control for cell viability, a similar test was carried out on plates carrying uracil (right panels).

domain (GAL4BD) and the corresponding wild-type partner to the activation domain (GAL4AD), or vice versa, obtaining in both cases similar results.

An interaction of P2α with the P1β mutants was detected in all cases, even with P1β5ntP2β, which is unable to bind to the ribosome (Figure 3A). In the P2β series, the wild-type P1α was able to interact only with P2βS2 and P2βD3, though to a reduced extent, especially in the first case. All the other tested proteins totally failed to allow cell growth and, therefore, seemed unable to interact with P1α (Figure 3B). An immunoblotting analysis confirmed that the mutant proteins fused to the Gal4 domain were present in the *S. cerevisiae* MaV203 cell extracts (Supplementary Data and Supplementary Figure S2).

A comparison of the two-hybrid and ribosome binding tests (Figures 2 and 3) indicated that there was not always a direct relationship between the capacity for ribosome binding and for P1/P2 association. Thus, protein P1β5ntP2β, which showed a practically normal level of association with P2α, was totally absent from the ribosome. In contrast, protein P2βS2S4, present in significant amounts in ribosomes from D56 and practically at the control level in D5, did not show interaction with P1α in the two-hybrid system. Since the P2β mutant result was totally unexpected, it was confirmed using affinity chromatography as an alternative experimental approach for testing interactions. C-terminal His6-tagged P1α was expressed in *S. cerevisiae* D5 together with P2βS2, P2βS2S4 or wild-type P2β. The supernatants (S100 fractions) from the cells were loaded in a Ni²⁺-NTA column, and the




Figure 4. Estimation of the interaction of P2β mutants with protein P1α using affinity chromatography. Supernatant S100 extracts (S) from *S. cerevisiae* cells expressing simultaneously the indicated P2β proteins and P1α-His6 were adsorbed in Ni²⁺-NTA resin. After washing, the fractions eluted with 100 mM imidazol were analyzed by SDS-PAGE and the P2β proteins detected by immunoblotting (E). A non-characterized protein detected by the antibody (N) can be used as loading control.

amount of each P2β specifically bound to resin was estimated by immunoblotting. As shown in Figure 4, while wild-type P2β is strongly retained by P1α-His6 in the column as expected, mutant P2βS2 is drastically reduced and mutant P2βS2S4 totally absent in the eluted fractions.

Modeling *S. cerevisiae* P1–P2 heterodimers and the (P0,P1α-P2β,P1β-P2α) pentamer

An accurate interpretation of the previous results would be notably helped by a high-resolution structure of the yeast stalk, which is not yet available. However, the recently reported structures of the human P2 homodimer (21), and the archaeal pentameric [P0(P1)₂(P1)₂(P1)₂] stalk complex (20), have allowed homology modeling of the structure of the equivalent yeast protein complexes. As expected, the two yeast heterodimer structures, P1α-P2β and P1β-P2α, are quite similar (Supplementary Figure S4A). They slightly diverge at the carboxyl terminal domains but the α-helix 1 nicely overlaps in both models. Since the P1 and P2 NTD sequence differ notably, the structure of the yeast heterodimers lacks the symmetry of the human P2 homodimer. Thus, the position of the P1 and P2 N-terminal in relation to the whole structure is significantly different (Supplementary Data and Supplementary Figure S4B).

The yeast stalk structure (Supplementary Figure S5) has been modeled taking into consideration two facts: (i) first, the P1α/P2β and P1β/P2α heterodimers bind to the first and second α-helix in the P0 C-terminal spine, respectively (17) and (ii) the data from the human stalk support that the P1 proteins from both heterodimers are probably facing each other in the pentamer (21). The modeled *S. cerevisiae* pentamer shows differences compared to the equivalent human structure mainly due to the disparity of the two yeast heterodimers. Among others, the angle formed by the two P0 α-helices is considerably more acute in yeast pentamer (Supplementary Data and Supplementary Figure S6).

DISCUSSION

Structural stability of the individual proteins

It was previously shown that the radical difference in protease sensitivity of P1 and P2 proteins lays in the first five amino acids of each protein (23). This report shows that just the mutation S2F totally abolishes the high protease sensitivity of protein P1 β . As previously reported (35,40), the presence in P1 β of a serine in the second position of the amino acid sequence induces removal of the starting Met and acetylation of the exposed NH₂-end. In contrast, the presence of a bulky Phe residue after the initiator Met blocked the N-amino end processing of proteins (Table 1). We have previously shown that N-terminal acetylation is not a degradation determinant for P proteins (23); therefore, cleavage of the first Met seems to be a requirement for degradation of P1 β , and probably for P1 proteins in general since all of them carry a small uncharged amino acid in the second position (41,42), which must induce N-amino end processing (43,44).

On the other hand, replacing the Lys2 in P2 β , an N-amino end processing blocker, by a serine has a limited effect on protein stability, in spite of the fact that the mutation induces extensive processing of the protein N-terminal. In contrast, mutation in position 3, which did not cause N-terminal processing (Table 1), results in a drastic decrease of protein P2 β stability either in the presence or in the absence of a serine in the second position.

These results indicate that the actual degradation determinants are different in both protein types. Thus, amino acids in positions 2 and 3 are determining the half-life of proteins P1 β and P2 β , respectively. For this reason, the same four initial amino acids, MKDL, which contains a P1 β stabilizing Lys in position 2 and a P2 β destabilizing Glu at position 3, can have an opposite effect in the two proteins, inducing stability of P1 β K2L4 and instability in P2 β D3. Probably, other elements may exist in the proteins, perhaps secondary or tertiary structural features, which together with the N-terminal determinants, define their different degradation sensitivity. Alternatively, the existence of different degradation mechanisms for each protein type cannot be excluded.

P1–P2 heterodimer formation

The most interesting observation regarding the capacity of the 12 kDa acidic stalk proteins to form P1–P2 heterodimers is the sensitivity of the process to mutations at the amino end as well as the different response of the two protein types to the modifications. While most P2 β mutants are unable to associate with their partner protein, similar modifications do not affect the P1 association capacity.

The association of two human P2 proteins to form a homodimer takes place mainly through symmetrical hydrophobic interactions between α -helices 1, 2 and 3 from both components (21). The modeled yeast heterodimer structures are more asymmetric and the N-ends of P1 and P2 proteins have a clearly different

structural environment. The P2 α -helix 1 is shorter than in P1, and its N-terminal is in a more internal position that provides more stabilizing interaction possibilities for the first amino acids (Supplementary Figure S4B) and might explain the different effects of similar mutations on both proteins. Thus, while in P1 β Ser2, which is actually the first residue due to the removal of the initiating methionine, and Ser4, can hardly be involved in important interactions, Met1 and Lys2 in P2 β can directly interact through hydrogen bonding with Asp15 and Leu11 lateral chains in P1 α α -helix 3 (Figure 5A). Residues Tyr3 or Leu4, whose mutation is required for total elimination of the P2 β associating capacity, are, however, oriented toward the inside and they cannot be directly involved in the interaction with P1 α but their substitution probably changes the protein conformation. Thus, Leu4 is part of the hydrophobic core formed by P2 β helices 1, 3 and 4, which can be seriously disturbed in the mutants. Taking into consideration all these facts, it is likely that the mutations in the P2 β N-end, in addition to directly affecting the interaction with P1 α , seem to alter the first turn of helix 1 inducing a conformational change that apparently affects the stability of the whole complex. The structural similarity of both yeast heterodimers as well as high sequence homology of the N-terminal domains of the proteins from both families make it highly probable that data obtained from P1 β and P2 β can be extended to P1 α and P2 α , respectively. In any case, confirmation of our structural predictions will require the crystal structure of the yeast stalk complex, which we hope will not take too long to be resolved.

As a whole, the data show the great structural relevance of the first turn of α -helix 1 in P2 proteins, but not in P1 proteins for heterodimer stability, and stress the differences in the structural role of the two proteins families have previously been reported using other experimental approaches (45).

Assembly into the yeast ribosomal stalk pentamer

In the modeled pentamer the main structural elements participating in the interaction with P0 are α -helices 1 and 4 from the N-terminal domain of the four acidic proteins (Supplementary Figure S5). Our results show that individually the first four amino acids of P2 β , which practically form the α -helix 1 first turn, partially contribute to the stability of the complex and all together are essential in maintaining the association of the heterodimer with stalk. In fact, the four amino acids show potential interactions with residues in the P0 α -helix as well as in P1 α -helix 1 and P2 β α -helix 3 (Figure 5B).

In contrast, modification of positions 2 and 4 in P1 β does not affect protein binding to the ribosome and apparently does not contribute much to the stability of the complex. Actually, in the modeled pentamer structure the first three residues of the protein are apparently not included in α -helix 1 (Supplementary Figure S5). A more extensive modification including the first five residues, which in this case affects helix 1 (Figure 5C), abolishes the protein capacity to bind to the ribosome. It seems, therefore, that as in the heterodimer formation, the




Figure 5. Close view of the amino ends from protein P2 β (A and B) and P1 β (A and C) in the modeled *S. cerevisiae* stalk heterodimer (A) and pentamer (B and C) shown in Supplementary Figures S4 and S5. (A) Protein P2 β α -helices 1, 2, 3 and 4 (cyan) and P1 α α -helix 3 (green) are shown. Lateral chains, names and sequence position of the first four amino acids from P2 β are shown (red) as well as those from the nearer amino acids in other P2 β regions (orange) and in P1 α α -helix 3 (magenta). (B) Helices from protein P2 β (blue), P1 α (magenta) and P0 (green) surrounding the first amino acids from P2 β α -helix 1 (red) are shown. The closest residues to these amino acids in the surrounding helices are marked in orange. (C) A closer view of the first amino acids (red) from protein P1 β (orange) is shown. The closer structural elements from proteins P2 α (cyan), P1 α (magenta) and P0 (green) are shown. The amino and carboxyl end of the P0 fragment are marked in panels. Note that the orientation of the model is opposite to that shown in Supplementary Figure S5.

initial amino acids of P1 and P2 proteins are not equally relevant.

The differences found in ribosome binding of P2 β mutants in *S. cerevisiae* D5 and D56 (Figure 2C) indicate that the presence of the protein P1 β , absent in the D56 cells, facilitates the association of the defective proteins to the ribosome. Protein P1 β can form P1 β /P2 α heterodimers that allow the assembly of a complete (P1 α /P2 β)–P0–(P1 β /P2 α) pentamer, which is missing in strain D56 and seems to stabilize the binding of the defective mutant P2 β –P1 α couples. These results support that the two heterodimers forming the ribosomal stalk are not independent as has been proposed for the human ribosomes (21).

Role of heterodimers in pentamer assembly

As commented on previously, P1 and P2 are able to form heterodimers in solution, and it is assumed that their interaction with the ribosome takes place in the associated state. If this is so, there must be a direct relationship between the protein capacity to form heterodimers and to bind to the ribosome. In general, our results are in

agreement with this statement but not in all cases. Some P2 β mutants are detected as bound to the ribosome in high amounts, but are seriously or even totally affected in their capacity to interact in solution with P1 α . Mutant P2 β S2S4 is probably the clearest example. This protein was unable to interact with P1 α and P2 α or even to form homo-dimers (Figure 3 and Supplementary Figure S3 in Supplementary Data) but was found in high proportions in the washed D56 and D5 ribosomes. Also, mutant P2 β S2 was present in the ribosomes of D5 and D56 in normal amounts but showed an important reduction in its capacity to interact directly with P1 α . It seems, therefore, that although wild-type P1 and P2 proteins are usually associated as highly stable P1/P2 heterodimers and interact with P0 as couples, a strong association is not an essential requirement for forming a functional stalk in the ribosome. The existence of weak associations, not detectable by standard techniques, which is stabilized upon binding to P0, cannot be excluded. Actually, the association of the acidic proteins with the ribosome, like the binding of P0 (46–48), might require the help of assembly factors, which can facilitate the process in the

case of acidic proteins which do not form stable couples in solution.

SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

ACKNOWLEDGEMENTS

We thank M.C. Fernández Moyano for expert technical assistance.

FUNDING

The Spanish Ministry of Science and Innovation (MICINN) (BFU2007-64280 to M.R. and BFU2009-09738 to J.P.G.B.); and Fundación Ramón Areces (Institutional Grant to C.B.M.S.O.). Funding for open access charge: Grants from Spanish Government.

Conflict of interest statement. None declared.

REFERENCES

- Gonzalo, P. and Reboud, J.P. (2003) The puzzling lateral flexible stalk of the ribosome. *Biol. Cell*, **95**, 179–193.
- Diaconu, M., Kothe, U., Schlunzen, F., Fischer, N., Harms, J.M., Tonevitsky, A.G., Stark, H., Rodnina, M.V. and Wahl, M.C. (2005) Structural basis for the function of the ribosomal L7/12 stalk in factor binding and GTPase activation. *Cell*, **121**, 991–1004.
- Ilag, L.L., Videler, H., McKay, A.R., Sobott, F., Fucini, P., Nierhaus, K.H. and Robinson, C.V. (2005) Heptameric (L12)6/L10 rather than canonical pentameric complexes are found by tandem MS of intact ribosomes from thermophilic bacteria. *Proc. Natl Acad. Sci. USA*, **102**, 8192–8197.
- Uchiyama, T. and Kominami, R. (1997) Binding of mammalian ribosomal protein complex P0–P1–P2 and protein L12 to the GTPase-associated domain of 28 S ribosomal RNA and effect on the accessibility to anti-28 S RNA autoantibody. *J. Biol. Chem.*, **272**, 3302–3308.
- Tchorzewski, M., Boldyreff, B., Issinger, O. and Grankowski, N. (2000) Analysis of the protein–protein interactions between the human acidic ribosomal P-proteins: evaluation by the two hybrid system. *Int. J. Biochem. Cell Biol.*, **32**, 737–746.
- Guarinos, E., Remacha, M. and Ballesta, J.P.G. (2001) Asymmetric interactions between the acidic P1 and P2 proteins in the *Saccharomyces cerevisiae* ribosomal stalk. *J. Biol. Chem.*, **276**, 32474–32479.
- Petersson, I., Hardy, S.J. and Liljas, A. (1976) The ribosomal protein L8 is a complex of L7/L12 and L10. *FEBS Lett.*, **64**, 135–138.
- Sanchez-Madrid, F., Reyes, R., Conde, P. and Ballesta, J.P.G. (1979) Acidic ribosomal proteins from eukaryotic cells. Effect on ribosomal functions. *Eur. J. Biochem.*, **98**, 409–416.
- Zinker, S. and Warner, J.R. (1976) The ribosomal proteins of *Saccharomyces cerevisiae*. Phosphorylated and exchangeable proteins. *J. Biol. Chem.*, **251**, 1799–1807.
- Tsurugi, K. and Ogata, K. (1985) Evidence for the exchangeability of acidic ribosomal proteins on cytoplasmic ribosomes in regenerating rat liver. *J. Biochem.*, **98**, 1427–1431.
- Scharf, K.-D. and Nover, L. (1987) Control of ribosome biosynthesis in plant cell cultures under heat shock conditions. II. Ribosomal proteins. *Biochim. Biophys. Acta*, **909**, 44–57.
- Remacha, M., Jimenez-Diaz, A., Santos, C., Zambrano, R., Briones, E., Rodriguez Gabriel, M.A., Guarinos, E. and Ballesta, J.P.G. (1995) The proteins P1, P2, and P0, components of the eukaryotic ribosome stalk. New structural and functional aspects. *Biochem. Cell Biol.*, **73**, 959–968.
- Saenz-Robles, M.T., Remacha, M., Vilella, M.D., Zinker, S. and Ballesta, J.P.G. (1990) The acidic ribosomal proteins as regulators of the eukaryotic ribosomal activity. *Biochim. Biophys. Acta*, **1050**, 51–55.
- Guarinos, E., Santos, C., Sanchez, A., Qiu, D.Y., Remacha, M. and Ballesta, J.P. (2003) Tag-mediated fractionation of yeast ribosome populations proves the monomeric organization of the eukaryotic ribosomal stalk structure. *Mol. Microbiol.*, **50**, 703–712.
- Remacha, M., Jimenez-Diaz, A., Bermejo, B., Rodriguez-Gabriel, M.A., Guarinos, E. and Ballesta, J.P.G. (1995) Ribosomal acidic phosphoproteins P1 and P2 are not required for cell viability but regulate the pattern of protein expression in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.*, **15**, 4754–4762.
- Perez-Fernandez, J., Remacha, M. and Ballesta, J.P. (2005) The acidic protein binding site is partially hidden in the free *Saccharomyces cerevisiae* ribosomal stalk protein P0. *Biochemistry*, **44**, 5532–5540.
- Krokowski, D., Boguszevska, A., Abramczyk, D., Liljas, A., Tchorzewski, M. and Grankowski, N. (2006) Yeast ribosomal P0 protein has two separate binding sites for P1/P2 proteins. *Mol. Microbiol.*, **60**, 386–400.
- Hagiya, A., Naganuma, T., Maki, Y., Ohta, J., Tohkairin, Y., Shimizu, T., Nomura, T., Hachimori, A. and Uchiyama, T. (2005) A mode of assembly of P0, P1, and P2 proteins at the GTPase-associated center in animal ribosome: *in vitro* analyses with P0 truncation mutants. *J. Biol. Chem.*, **280**, 39193–39199.
- Briceno, V., Camargo, H., Remacha, M., Santos, C. and Ballesta, J.P. (2008) Structural and functional characterization of the amino terminal domain of the yeast ribosomal stalk P1 and P2 proteins. *Int. J. Biochem. Cell Biol.*, **41**, 1315–1322.
- Naganuma, T., Nomura, N., Yao, M., Mochizuki, M., Uchiyama, T. and Tanaka, I. (2010) Structural basis for translation factor recruitment to the eukaryotic/archaeal ribosomes. *J. Biol. Chem.*, **285**, 4747–4756.
- Lee, K.M., Yu, C.W., Chan, D.S., Chiu, T.Y., Zhu, G., Sze, K.H., Shaw, P.C. and Wong, K.B. (2010) Solution structure of the dimerization domain of ribosomal protein P2 provides insights for the structural organization of eukaryotic stalk. *Nucleic Acids Res.*, **38**, 5206–5215.
- Bermejo, B., Remacha, M., Ortiz-Reyes, B., Santos, C. and Ballesta, J.P. (1994) Effect of acidic ribosomal phosphoprotein mRNA 5'-untranslated region on gene expression and protein accumulation. *J. Biol. Chem.*, **269**, 3968–3975.
- Nussbaumer, G., Remacha, M. and Ballesta, J.P. (2000) Phosphorylation and N-terminal region of yeast ribosomal protein P1 mediate its degradation, which is prevented by protein P2. *EMBO J.*, **19**, 6075–6084.
- Remacha, M., Santos, C., Bermejo, B., Naranda, T. and Ballesta, J.P.G. (1992) Stable binding of the eukaryotic acidic phosphoproteins to the ribosome is not an absolute requirement for *in vivo* protein synthesis. *J. Biol. Chem.*, **267**, 12061–12067.
- Remacha, M., Santos, C. and Ballesta, J.P. (1990) Disruption of single-copy genes encoding acidic ribosomal proteins in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.*, **10**, 2182–2190.
- Naranda, T. and Ballesta, J.P.G. (1991) Phosphorylation of acidic proteins controls the activity of the ribosome. *Proc. Natl Acad. Sci. USA*, **88**, 10563–10567.
- Francisco-Velilla, R. and Remacha, M. (2010) *In vivo* formation of a stable pentameric (P2alpha/P1beta)-P0-(P1alpha/P2beta) ribosomal stalk complex in *Saccharomyces cerevisiae*. *Yeast*, **9**, 693–704.
- Sambrook, J. and Russell, D.W. (2006) *Molecular Cloning: A Laboratory Manual*, 3rd edn. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Dieffenbach, C.W. and Dveksler, G.S. (1995) *PCR Primer. A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Zambrano, R., Briones, E., Remacha, M. and Ballesta, J.P.G. (1997) Phosphorylation of the acidic ribosomal P proteins in *Saccharomyces cerevisiae*. A reappraisal. *Biochemistry*, **36**, 14439–14446.
- Vilella, M.D., Remacha, M., Ortiz, B.L., Mendez, E. and Ballesta, J.P.G. (1991) Characterization of the yeast acidic ribosomal phosphoproteins using monoclonal antibodies.

- Proteins L44/L45 and L44' have different functional roles. *Eur. J. Biochem.*, **196**, 407–414.
32. Vidal, M., Brachmann, R.K., Fattaey, A., Harlow, E. and Boeke, J.D. (1996) Reverse two-hybrid and one-hybrid systems to detect dissociation of protein–protein and DNA–protein interactions. *Proc. Natl Acad. Sci. USA*, **93**, 10315–10320.
 33. Eswar, N., Webb, B., Marti-Renom, M.A., Madhusudhan, M.S., Eramian, D., Shen, M.Y., Pieper, U. and Sali, A. (2006) Comparative protein structure modeling using Modeller. *Curr. Protoc. Bioinform.*, **Chapter 5**, Unit 5.6.
 34. Phillips, J.C., Braun, R., Wang, W., Gumbart, J., Tajkhorshid, E., Villa, E., Chipot, C., Skeel, R.D., Kalé, L. and Schulten, K. (2005) Scalable molecular dynamics with NAMD. *J. Computat. Chem.*, **26**, 1781–1802.
 35. Santos, C., Ortiz-Reyes, B.L., Naranda, T., Remacha, M. and Ballesta, J.P.G. (1993) The acidic phosphoproteins from *Saccharomyces cerevisiae* ribosomes. NH₂-terminal acetylation is a conserved difference between P1 and P2 proteins. *Biochemistry*, **32**, 4231–4236.
 36. Bachmair, A., Finley, D. and Varshavsky, A. (1986) *In vivo* half-life of a protein is a function of its amino-terminal residue. *Science*, **234**, 179–186.
 37. Mayer, A., Siegel, N.R., Schwartz, A.L. and Ciechanover, A. (1989) Degradation of proteins with acetylated amino termini by the ubiquitin system. *Science*, **244**, 1480–1483.
 38. Naranda, T., Remacha, M. and Ballesta, J.P. (1993) The activity-controlling phosphorylation site is not the same in the four acidic ribosomal proteins from *Saccharomyces cerevisiae*. *J. Biol. Chem.*, **268**, 2451–2457.
 39. Tchorzewski, M. (2002) The acidic ribosomal P proteins. *Int. J. Biochem. Cell Biol.*, **34**, 911–915.
 40. Takakura, H., Tsunasawa, S., Miyagi, M. and Warner, J.R. (1992) NH₂-terminal acetylation of ribosomal proteins of *Saccharomyces cerevisiae*. *J. Biol. Chem.*, **267**, 5442–5445.
 41. Ballesta, J.P.G. and Remacha, M. (1996) The large ribosomal subunit stalk as a regulatory element of the eukaryotic translational machinery. *Progr. Nucleic Acid. Res. Mol. Biol.*, **55**, 157–193.
 42. Grela, P., Bernado, P., Svergun, D., Kwiatowski, J., Abramczyk, D., Grankowski, N. and Tchorzewski, M. (2008) Structural relationships among the ribosomal stalk proteins from the three domains of life. *J. Mol. Evol.*, **67**, 154–167.
 43. Boissel, J.P., Kasper, T.J. and Bunn, H.F. (1988) Cotranslational amino-terminal processing of cytosolic proteins. Cell-free expression of site-directed mutants of human hemoglobin. *J. Biol. Chem.*, **263**, 8443–8449.
 44. Hirel, P.H., Schmitter, M.J., Dessen, P., Fayat, G. and Blanquet, S. (1989) Extent of N-terminal methionine excision from *Escherichia coli* proteins is governed by the side-chain length of the penultimate amino acid. *Proc. Natl Acad. Sci. USA*, **86**, 8247–8251.
 45. Qiu, D., Parada, P., Marcos, A.G., Cardenas, D., Remacha, M. and Ballesta, J.P. (2006) Different roles of P1 and P2 *Saccharomyces cerevisiae* ribosomal stalk proteins revealed by cross-linking. *Mol. Microbiol.*, **62**, 1191–1202.
 46. Lo, K.Y., Li, Z., Wang, F., Marcotte, E.M. and Johnson, A.W. (2009) Ribosome stalk assembly requires the dual-specificity phosphatase Yvh1 for the exchange of Mrt4 with P0. *J. Cell. Biol.*, **186**, 849–862.
 47. Kemmler, S., Occhipinti, L., Veisu, M. and Panse, V.G. (2009) Yvh1 is required for a late maturation step in the 60S biogenesis pathway. *J. Cell. Biol.*, **186**, 863–880.
 48. Rodriguez-Mateos, M., Garcia-Gomez, J.J., Francisco-Velilla, R., Remacha, M., de la Cruz, J. and Ballesta, J.P. (2009) Role and dynamics of the ribosomal protein P0 and its related trans-acting factor Mrt4 during ribosome assembly in *Saccharomyces cerevisiae*. *Nucleic Acids Res.*, **37**, 7519–7532.